




LOW STARCH

PELLET

50 LBS
(22.7 KG)



IDEAL FOR:

Mature horses, ponies, metabolic horses



FEATURES:

Whole grain, alfalfa & molasses free, low starch & sugar, lower potassium



HELPS SUPPORT:

Metabolic disease, HYPP, allergies

Triple Crown Low Starch contains no whole grains, molasses or alfalfa meal, ingredients many horse owners associate with hyperactivity and allergies. With fewer carbohydrates than most hay and pasture, this feed provides additional calories for performance horses and horses with elevated levels of metabolism. By utilizing high quality fiber sources and added fat, including rice bran and flaxseed, essential calories are provided while keeping sugar and starch levels low.

PROTEIN 13% • FAT 6% • FIBER 18% • 1,428 KCAL/LB

GUARANTEED ANALYSIS

Crude Protein (min.)	13.00%
Lysine (min.)	0.75%
Methionine (min.)	0.25%
Threonine (min.)	0.70%
Tryptophan (min.)	0.15%
Leucine (min.)	0.90%
Crude Fat (min.)	6.00%
Crude Fiber (max.)	18.00%
ADF (max.)	22.00%
NDF (max.)	44.00%
Calcium (min.)	0.75%
Calcium (max.)	1.25%
Phosphorus (min.)	0.60%
Magnesium (min.)	0.50%
Potassium (min.)	0.75%
Sodium (min.)	0.50%
Sodium (max.)	1.00%
Salt (min.)	0.75%
Salt (max.)	1.50%
Omega-3 Fatty Acids (min.)	0.60%
Omega-6 Fatty Acids (min.)	2.90%
Selenium (min.)	0.60 ppm
Selenium (max.)	1.60 ppm
Zinc (min.)	220 ppm
Manganese (min.)	100ppm
Cobalt (min.)	0.20 ppm
Copper (min.)	50 ppm
Iron (min.)	100 ppm
Vitamin A (min.)	6,000 IU/lb.
Vitamin D (min.)	1,200 IU/lb.
Vitamin E (min.)	205 IU/lb.
Riboflavin-B2 (min.)	3.20 mg/lb.
Thiamin-B1 (min.)	8.00 mg/lb.
Ascorbic Acid (min.)	55 mg/lb.
Biotin (min.)	0.25 mg/lb.
Total Direct Fed Microbials (min.)	3.64 billion CFU/lb.
Lactobacillus Acidophilus Fermentation Product (min.)	908 million CFU/lb.
Enterococcus Faecium Fermentation Product (min.)	908 million CFU/lb.
Bacillus Subtilis Dried Fermentation Product (min.)	25 million CFU/lb.
Saccharomyces Cerevisiae Yeast Culture (min.)	1.80 billion CFU/lb.
Cellulase ¹ (Trichoderma Longibrachiatum Fermentation Extract) (min.)	0.68 million CFU/lb.
Protease ² (Bacillus Subtilis Fermentation Extract) (min.)	1,430 HUT/lb.

INGREDIENTS

(Free from Restricted Ruminant Protein Products per Title 21, CFR 589.2000/2001) Wheat Middlings, Soybean Hulls, Ground Beet Pulp, Distillers Dried Grains, Soybean Oil, Ground Flaxseed, Stabilized Rice Bran, Calcium Carbonate, Salt, Sodium Bicarbonate, Dicalcium Phosphate, Yeast Extract, Yeast Culture, Calcium Silicate, Dried Enterococcus Faecium Fermentation Product, Dried Lactobacillus Acidophilus Fermentation Product, Dried Bacillus Subtilis Fermentation Product, Dried Trichoderma Longibrachiatum Fermentation Extract, Dried Bacillus Subtilis Fermentation Extract, Zinc Hydroxy Chloride, Manganese Hydroxy Chloride, Copper Hydroxy Chloride, Vegetable Oil, Butyric Acid, Zinc Oxide, Peppermint Essential Oil, Kelp Meal, Magnesium Oxide, Fenugreek Seed, Lecithin, Anise, Magnesium Proteinates, Selenium Yeast, L-Lysine, DL-methionine, L-Threonine, L-Leucine, Ascorbic Acid (Source of Vitamin C), Niacin Supplement, Biotin, Vitamin A Supplement, Vitamin E Supplement, Thiamine Mononitrate, Beta Carotene, Calcium Pantothenate, Riboflavin Supplement, Pyridoxine Hydrochloride, Vitamin B12 Supplement, Vitamin D3 Supplement, Choline Chloride, Menadione Sodium Bisulfite Complex (Source of Vitamin K Activity), Folic Acid, Hydrated Sodium Calcium Aluminosilicate, Manganese Sulfate, Lignin Sulfonate, Brewers Dried Yeast, Zinc Sulfate, Copper Sulfate, Cobalt Proteinates, Iron Amino Acid Complex, Ethylenediamine Dihydroiodide, (Propionic Acid, Sodium Benzoate, Potassium Sorbate (Preservatives)).

FEEDING DIRECTIONS

MATURE HORSES

900 – 1,100 lb. 6 lb. per day

PONIES

500 – 600 lb. 4 lb. per day

OTHER CONSIDERATIONS

If feeding less than 6 lb. to a 900 – 1,100 lb. horse, supplement with 1 lb. per day of Triple Crown Balancer or Triple Crown Balancer Gold in order to meet recommended levels of vitamins, minerals and digestive support.

Triple Crown Low Starch is a complete diet and can replace all or part of the fiber portion of your horse's diet. If feeding as the entire ration and replacing all the hay in the diet, feed 1.0 to 2% of body weight daily. Triple Crown Low Starch can be soaked with warm water for horses with poor teeth, respiratory problems or for horses prone to choke.

Adjust the amount of Triple Crown Low Starch up or down as needed to maintain desired body condition. Do not feed more than 5 lb. at a single meal. Allow 3 to 4 hours between meals.

1. One cellulase unit (CU) is defined as the amount of activity that will produce a relative fluidity change of 1 (determined with a calibrated viscometer) in 5 minutes in a defined carboxymethylcellulose substrate at pH 4.5 and 40°C.
2. One hemoglobin unit (HUT) of proteolytic (protease) activity is defined as that amount of enzyme that produces, in one minute under the specified conditions, a hydrolysate whose absorbance at 275nm is the same as that of a solution containing 1.10g per mL of tyrosine in 0.006N hydrochloric acid. Estimated values determined by Equi-Analytical, Ithaca, NY.